

IV071000 **Double Spike Adaptor**

INSTRUCTION FOR USE

DESCRIPTION

The m|devices Double Spike Adaptor is designed with two spikes, two roller regulators and a septum connector.

The inner tubing diameter is 3mm and total length is 30cm.

FOR USE BY A QUALIFIED CLINICIAN. THE BELOW IS ONLY A SUGGESTION AND FACILITY PROTOCOL MUST BE FOLLOWED FOR ALL CLINICAL PROCEDURES WHERE THIS PRODUCT IS USED.

CAUTION

- · Single use
- DO NOT re-sterilise.
- DO NOT store at extreme temperatures and humidity, avoid direct sunlight.
- STERILE (EO), DO NOT use if the package or product has been damaged or contaminated
- . EU Notice: any serious incident that has occurred in relation to the device should be reported to the manufacture and the competent authority of the Member State in which the user and /or patient is established.

STEPS

- 1. Remove product from packaging, maintaining aseptic technique.
- 2. Close roller regulators on the double spike adaptor.
- 3. Remove protective cap from spikes and insert into fluid bag/containers.
- 4. Suspend bag/containers.
- 5. Close all roller regulators and/or click clamps on the administration set that will be connected to the double spike set.
- 6. Insert the administration set spike into the in-line port of the double spike adapter until the connection is secure.
- 7. Open roller regulators on the double spike adaptor to allow the fluid to prime to the inline port through to the IV administration set.
- 8. To further prime past the in-line port, follow instructions to the administration set until all air has been expelled from the line to the luer adaptor.

PRECAUTION

• The security of all connections should be checked once established and monitored during use to prevent disconnection.



mldevices

27 Llewellyn Avenue Villawood NSW 2163 Australia T +61(0)2 8718 2800 F +61(0)2 8718 2801



BQ Plus Medical Co., Ltd No. 18. Che Ye Road Che Dun Town, Songjiang 201611 Shanghai, China



Brehmstr. 56 40239 Duesseldorf Germany









